

Original Article

## POLLUTION ASSESSMENT: THE INFLUENCE OF MUNICIPAL SEWAGE ON WATER BODIES IN OBIO/AKPOR

*Ifeanyi Chukwuemeka Nwankwo and Abdulrazaq Temitope Oladimeji*

Department of Microbiology,  
Rivers State University of Science and  
Technology,  
Port Harcourt, Rivers State, Nigeria  
DOI:<https://doi.org/10.5281/zenodo.16024365>

**Abstract:** This study evaluates the impact of municipal sewage disposal on the quality of surface water in the Obio/Akpor Local Government Area of Rivers State, Nigeria. Water samples were collected from five different stations along the Rumuolemini community and adjoining creeks along the Iwofe Road axis over a six-month period (June–November). Sampling was conducted three times weekly using standard analytical methods. Heavy metals including zinc (Zn), manganese (Mn), iron (Fe), copper (Cu), and lead (Pb) were quantified using Atomic Absorption Spectrophotometry (AAS), while microbial quality was assessed using the multiple tube fermentation technique to determine the Most Probable Number (MPN) of coliform bacteria.

Results indicated significantly high microbial loads across the stations, with Station 1 showing the highest faecal and total coliform counts at  $25,691.00 \pm 1.41$  and  $35,602.5 \pm 3.54$  MPN/100 mL, respectively. Total heterotrophic bacterial count peaked at Station 4 ( $6.10 \pm 0.14$  log CFU/mL). Significant variations in coliform levels were observed among the stations, particularly for total coliforms. Heavy metal concentrations were ranked in descending order as  $Fe > Mn > Cu > Zn$ , with iron levels exceeding acceptable thresholds in several samples.

The elevated levels of faecal contamination and heavy metals suggest that the receiving water bodies are significantly polluted due to untreated sewage discharge. These conditions pose serious risks to human and aquatic health, potentially leading to outbreaks of waterborne diseases such as typhoid fever, cholera, dysentery, and gastroenteritis. Furthermore, high iron content may contribute to water discoloration and habitat degradation.

The study concludes that the current practice of discharging untreated municipal wastewater into local rivers compromises water safety and quality. It recommends the implementation of effective wastewater treatment techniques such as phytoremediation to reduce pollutant loads. Additionally, treated effluents could be repurposed for agricultural irrigation rather than released directly into the environment. Urgent regulatory and infrastructural interventions are required to restore ecological balance and safeguard public health in affected communities.

**Keywords:** sewage disposal, water quality, indicator microorganisms, heavy metals, pollution, diseases, Rumuolumeni

### INTRODUCTION

Rivers State and its territory is bounded with waters and streams yet there is no potable water supply due to various industrial and anthropogenic activities within the urban centres of the State. Sources of drinking water could get contaminated by sewage in various ways such as from septic tanks, leaking sewer lines, sludge used for land application, and from partially treated wastewater. This could bring about anaerobic conditions which would adversely affect aquatic life (Spellman and Drinan, 2000), including fish stocks, and the suffering of the

## **Original Article**

human diet. Also, it would cause appreciable health risks as water laden with untreated sewage is a potential source of water-borne disease-causing organisms. The effects can be far-reaching if such water is used to grow crops that are then eaten uncooked; in which case diseases can spread to whole communities.

It is well known that sewage and septic tank effluents are sources of the causative agents of gastrointestinal ailments such as diarrhea, dysentery, typhoid and cholera (Van and Pur, 1990; Bicki, 2001; Burabai *et al.*, 2007; Ochuko and Thaddeus, 2013; Shafi *et al.*, 2013; Ogbonna, 2014). In Nigeria, such ailments have been traced to the consumption of water polluted by untreated waste (UNEP, 2005; Ochuko and Thaddeus, 2013).

One of the major concerns in the Niger Delta is that the soils in the coastal plains are characterized by their hydraulically restrictive layers and seasonally high water table levels (Daniels and Weaver, 1987; Burabai *et al.*, 2007). If there are leakages from septic tanks it becomes easy for water to seep into the land surface and run off into surface water or flow directly into the water table to cause contamination. Also, the seasonal high water table conditions help porous soil structures to allow raw sewage to empty into the groundwater table. This would make percolation impossible since the soil is saturated, leading to infiltration and ex-filtration process in the septic tank (Daniels and Weaver, 1987). Effluents arising from such indiscriminate disposal could adversely affect the quality of soil and water because such processes generate and release enormous waste containing nitrogenous compounds into the terrestrial and aquatic environments or habitats. Thus the accumulation of the waste materials may result in toxic and hazardous effect on both humans and other inhabitants of the environment (Stewart, 2005). Such impact or damage may diminish the local biodiversity and reduce the possibility for many species to find shelter and feeding grounds.

Organic pollutants such as domestic sewage, urban run-off, farm wastes and effluents from food processing industries, brewing industries, dairies, abattoirs, tanneries, textile and paper making factories have diverse adverse effects. These pollutants are biodegradable and are easily oxidized by making use of the dissolved oxygen in water. However, as dissolved oxygen drops, fish and other aquatic life are threatened or killed in extreme cases. The number of algae and bacteria is increased; macrophyte levels are also adversely affected due to light reduction rendering the bed of the river unstable for plants (Oribhor, 2016). This study was therefore undertaken to assess the impact of municipal sewage disposal in water bodies around Obio/Akpor Local Government Area of Rivers State and its environs. It is expected that the study would help to address potential short- and long-term water quality issues from the various activities associated with the municipal sewage disposal which had hitherto threatened the integrity of many rivers and coastlines in the area.

## **Materials and Methods**

The sampling was carried out between June and November 2014. The locations of the sampling sites were established using a Garmin 45 Geographical Positioning System (GPS) (Fig 1).

Original Article




Fig 1. Map of Port Harcourt Metropolis showing sampling areas

Water samples were collected from the selected sites every month for six months from the Iwofe creek of Rumuolumini River. The samples were collected in a 1-litre plastic (1000 ml) bottles that had previously been carefully cleaned and rinsed thoroughly with distilled water (APHA, 1998).

Water samples for heavy metals were collected in a plastic bottle that had previously been soaked in concentrated nitric acid for 24 hours and rinsed with de-ionised water and brought to a pH of  $\leq 2$  with concentrated  $H_2SO_4$  or  $HNO_3$  and then stored in ice prior to laboratory analysis (Barley and Gardner, 1977). At the laboratory, the samples were stored in refrigerators at  $4^\circ C$  until analysis.



**Original Article**



**Fig 2: Sewage discharge point at Iwofe**



**Fig 3: Aquatic Microphytes on the water body Coliform count**

Water quality was determined by the standard multiple tube fermentation technique. Coliforms were tested for by inoculation of samples into tubes of lactose broth. The threetube procedure using lactose broth (Collins and Lyne, 1976; Bakare *et al.*, 2003) was used to the most probable number (MPN) of coliform bacilli (Collins and

## Original Article

Lyne, 1976). Water samples (10 ml, 1 ml and 0.1 ml) were inoculated into tubes with 10 ml of lactose broth and incubated at  $35 \pm 0.5^\circ\text{C}$ . Gas formation within 24 h of inoculation indicated positive presumptive test. Cultures showing no production of gas in 48h were considered negative. Samples from tubes showing gas were inoculated on to Eosine- methylene-blue agar; and one or more typical colonies were picked off into Brilliant Green Bile broth (Coyne and Howell, 1994) and studied microscopically for organisms with morphological and staining properties of coliform bacilli.

### Heavy metal analysis

Fifty millilitres of water samples in duplicate were digested with 10 ml of concentrated nitric acid in a pyrex conical flask. The solution, after digestion was cooled, filtered through a Whatman filter paper No. 42 into a 100 ml volumetric flask and made up to 100 ml mark with de-ionised water. A blank was also prepared in the same manner and method using deionised water so as to check the accuracy of the method. Stock standards (100 mg/l) for the various metals were prepared using standard (BDH, Analar grade) and the concentrated 30% nitric acid. The filtrates were analysed for heavy metals using Atomic Absorption Spectrophotometer (Model Perkin-Elmer 31000) and the calibration curve obtained for five dilutions preparation from the standards. The numerous data generated were subjected to basic statistical approach. The descriptive statistics were calculated using Microsoft Excel XP for Windows. The statistical analysis was to determine whether the principal component or heavy metal variables fall into different groups with possible same source(s) of entry into aquatic ecosystem.

## RESULTS

### Microbial counts

The microbial counts from the surface water bodies where the raw sewage was discharged was found to be above the threshold level, however the counts varied from station to station where sampling was carried out (Table 1). Faecal coliforms occurred more in station 1 where the raw municipal sewage were discharged, with a mean count of  $25691.00 \pm 1.41$  MPN index/100 ml. Similarly, total coliforms had the highest proportion of  $35602.00 \pm 3.54$  MPN index/100 ml in station 1 while total heterotrophic bacterial counts were  $6.10 \pm 0.14 \times 10^7$  cfu/ml in station 4. This trend was also observed in stations 2 and 3 for faecal coliforms which had  $19749.00 \pm 281.43$  MPN index/100 ml and  $29451.00 \pm 1.41$  MPN index/100ml for total coliforms respectively while station 5 had  $4.9 \pm 0.21 \times 10^7$  cfu/ml for total heterotrophic bacterial count. The occurrence of these indicator organisms was highly variable with significant differences between the raw sewage delivery point and other sampling sites along the zones examined.

**Table 1: Microbial counts in surface water at study area**

Stations	Faecal Coliforms (MPN Index/100ml)	Total Coliforms (MPN index/100ml)	Total Heterotrophic Bacteria ( $10^7$ cu/ml)
Station 1	$25691.00 \pm 1.41^c$	$35602.5 \pm 3.54^e$	$3.45 \pm 0.07^a$
Station 2	$19749.00 \pm 281.43^b$	$23665.00 \pm 134.35^c$	$3.20 \pm 0.00^a$

## Original Article

<b>Station 3</b>	13190.00±707.11 <sup>a</sup>	29451.00±1.41 <sup>d</sup>	3.35±0.07 <sup>a</sup>
<b>Station 4</b>	13861.00±1.41 <sup>a</sup>	16405.00±7.07 <sup>b</sup>	6.10±0.14 <sup>c</sup>

<b>Station 5</b>	13210.00±14.14 <sup>a</sup>	14805.00±70.07 <sup>a</sup>	4.9±0.21 <sup>b</sup>
------------------	-----------------------------	-----------------------------	-----------------------

Means in the same column with the same letter are not significantly different at 5% level of significance according to LSD test.

**Key:** Station 1: Sewage discharge point; Station 2: Creek junction; Station 3: Eagle Cement River; Station 4: Eagle Island Creek located between the Agip and Slaughter; Station 5: Ntanwogba Creek by Cherubim Road, Mile 3

Furthermore, there was no significant difference ( $p > 0.05$ ) in the occurrence of faecal coliforms between stations 3, 4 and 5 while there was a significant difference ( $p > 0.05$ ) in the number of total coliforms between all stations sampled. Also, there was a 0.05 level of significance in the total heterotrophic bacterial counts from stations 4 and 5 while there was no difference in the counts for stations 1, 2 and 3.

### Heavy metal levels

The mean concentrations of the heavy metals in the creek of Rumuolumini community are shown in Table 2. The values for iron content ranged from 0.31±0.002-13.14±0.014 ppm across the stations but were more in stations 1 and 2 with a total of 12.5±0.7 and 13.14±0.014 ppm respectively. Zinc, Copper and Manganese were detected from the samples. Lead was not detected in all the samples. The values for zinc were significantly different in trace amounts at 0.05 levels among the stations while managanese were not significantly different from stations 1, 2 and 3 but varied in concentrations in stations 4 and 5.

**Table 2: Heavy Metal Concentrations in surface water at the study areas**

Stations	Zinc (ppm)	Copper (ppm)	Lead (ppm)	Iron (ppm)	Manganese (ppm)
<b>Station 1</b>	0.41±0.003 <sup>e</sup>	0.16±0.01 <sup>d</sup>	0.00±0.00 <sup>a</sup>	12.5±0.7 <sup>c</sup>	0.35±0.07 <sup>b</sup>
<b>Station 2</b>	0.18±0.01 <sup>c</sup>	0.10±0.00 <sup>c</sup>	0.00±0.00 <sup>a</sup>	13.14±0.014 <sup>c</sup>	0.24±0.001 <sup>b</sup>
<b>Station 3</b>	0.34±0.01 <sup>d</sup>	0.08±0.00 <sup>b</sup>	0.00±0.00 <sup>a</sup>	0.41±0.001 <sup>a</sup>	0.32±0.005 <sup>b</sup>
<b>Station 4</b>	0.15±0.001 <sup>b</sup>	0.00±0.00 <sup>a</sup>	0.00±0.00 <sup>a</sup>	2.42±0.28 <sup>b</sup>	0.11±0.001 <sup>a</sup>
<b>Station 5</b>	0.05±0.01 <sup>a</sup>	0.00±0.00 <sup>a</sup>	0.00±0.00 <sup>a</sup>	0.31±0.002 <sup>a</sup>	0.08±0.00 <sup>a</sup>

Means in the same column with the same letter are not significantly different at 5% level of significance according to LSD test.

## DISCUSSION

### Microbial Counts

The water body in the vicinity where the municipal sewage wastes were discharged or used as dumpsites showed obvious deterioration in quality based on the number of faecal coliforms obtained. The MPN index observed for the water samples showed that the faecal coliform counts were above the permissible limit of WHO (2003, 2009), indicating gross pollution of the water body and its transition towards eutrophic state. According to Feng *et al* (2007), faecal coliforms are found in the intestinal canal of man and warm-blooded animals and are usually discharged alongside with faeces. The high counts of faecal coliforms are attributable to indiscriminate disposal of the municipal sewage along the banks of water bodies. Sometimes it may also be due to run-offs from urban solid waste disposal sites located around such vicinities which contain domestic animal and human faecal materials (Chapman, 1996; Agbolade *et al.*, 2009; 2010; Sharma *et al.*, 2010; Sadat *et al.*, 2011; Shafi *et al.*, 2013). Due to changes in agricultural methods, intensive farming operations utilize increased use of animal wastes as organic manure in most agricultural fields which also results in an increased pollution of rivers and streams especially through runoffs of leachates which drain into the water bodies (Gelt,



## **Original Article**

1998). This situation may cause contamination and transmission of diseases through water and food contaminated with human waste. Even though the people do not depend solely on these water bodies as their sources of water supply, the spate of water shortages could turn the tide as challenge. The high proliferation of heterotrophic bacteria in all the stations studied may have posed a challenge for competition amongst other groups of bacteria that could not survive for a long time. This might be as a result of increased microbial load washed into the river from the dumpsite by rain and the fact that more nutrients are brought in through leaching and run offs which eventually settle at the bottom of the river, leading to increased nutrient levels which encouraged rapid multiplication of bacteria (Adesemoye *et al.*, 2006).

### **Heavy metal concentrations**

The results of the mean concentrations of the heavy metals determined in the water body show that Fe, Zn, Mn and Cu were detected in various amounts with iron content exceeding tolerable limits (Table 2). Such amounts are a threat to river ecosystems due to increased turbidity resulting in accelerated sedimentation rates which changes the nature of water, thus affecting biological communities and physical habitats within the water body. The higher values of Fe and Zn could possibly be as a result of anthropogenic input vis-à-vis urbanization due to wastes from industrial and household materials, storm water runoff, precipitation carrying zinc from wear and tear of car tyres, corrosion of galvanized steel structures and leachates from refuse dumps carried into such surface water bodies. Such changes in sediment regime may also be caused by both land-based (such as catchments development, production-forest harvesting, road building) and water-based (bridge construction, eradication of noxious vegetation) activities (Nartey *et al.*, 2012 Seiyaboh *et al.*, 2013) which therefore defaces the aesthetic effect of affected water bodies.

Manganese occurs in surface waters that are low in oxygen and often does so with iron. When oxidized in aerobic waters, the oxide builds up in distribution causing severe discoloration at concentrations above 50.0 µg/l (WHO, 2009). The menace of water hyacinth or macrophytes in the rivers and streams is to some extent, attributed to the unintended fertilization of aquatic ecosystems by run-off from the sewage discharges onto such water bodies. The presence of manganese may be due to discharges from industrial facilities or due to indiscriminate sewage disposal or leachate from waste dumpsites around such areas (USEPA, 2009). Because the communities where these activities are undergoing various industrial activities like the cement industry and platforms for oil lifting activities.

### **CONCLUSION**

The present study was focused on the effect of municipal sewage disposal in some parts of Port Harcourt Metropolis. Monitoring these indicator organisms is an easy way to assess health risks due to bacterial contamination of surface waters. The presence of these physiologic groups in the samples is an indication of faecal contamination of the water bodies (Prescott *et al.*, 2005). This is possible since the sewage dumpsite is discharged within and around the water body. Through surface run-off, some of the faecal materials are carried to the nearby water body, leading to the presence of the coliform bacteria. Untreated faecal materials that contain faecal coliforms add excess organic material to the water. The decay of these materials depletes the water of oxygen which may result in killing of fishes and other aquatic life (Spellman and Drinan, 2000; Nartey *et al.*, 2012). According to UNICEF (1999), if water is found to contain faecal indicator bacteria, it is considered unsafe for human consumption.

The result of this work has clearly shown that the water samples from different places of Obio/Akpor LGA has become grossly polluted and anything affecting this habitat (water body) also affects human food, health and general well-being. This could also affect the health of the people that are constantly exposed to it or use the water contaminated by sewage. This work is very useful for efficient pollution supervision and for drawing pollution control strategy for water and soils within Obio/ Akpor LGA in Rivers state. This study also calls for the management of waste generated that may pose health associated risks and hazards.

## **Original Article**

### **References**

- Adesemoye, A O; Opere, B O and Makinde, S C O (2006). Microbial content of abattoir waste water and its contaminated soil in Lagos, Nigeria. *African Journal of Biotechnology* 5(20): 1963-1968.
- Adeyemo, O K; Ayodeji, I O and Aiki-raji, C O (2002): The water quality and the sanitary conditions in a major abattoir (Bodija) in Ibadan. *African Journal of Biomedical Research*, 5: 51-55
- Agbolade, O M; Oni, T T; Fagunwa, O E; Lawal, K M and Adesemowo, A (2009). Fecal contamination of dumpsites in some communities in Ijebu-North, South-Western Nigeria. *Nigeria Journal of Parasitology* 30(2):57-60
- Agbolade, O M; Adesanya, O O; Olayiwola, T O and Agu, G C (2010). Fecal and heavy metal contamination of some freshwaters and their vicinities in Ijebu-North, Southwestern Nigeria. *Journal of Toxicology and Environmental Health Sciences* 2(3): 32-37
- Alonge, D O (1991). *Textbook of Meat Hygiene in the Tropics* Farmcoe Press, Ibadan 58pp
- American Public Health Association, APHA) (1995). "Standard Methods for the Examination of Water and Wastewater. 20th Edition, American Public Health Association (APHA), American Water Works Association, (AWWA), Water Environment Federation (WEF)," Washington DC.
- American Public Health Association, APHA (1998). *Standard Methods for Examination of Water and Wastewater*. 20th edition. American Public Health Association, Washington D.C.
- Bakare, A A, Lateef A, Amuda O S and Afolabi R O (2003). The Aquatic toxicity and characterization of chemical and microbiological constituents of water samples from Oba River, Odo-oba, Nigeria. *Asian Journal of Microbiology Biotechnology and Environmental Science* 5:11-17.
- Barley, G E and Gardner, D (1977). Sampling and Storage of natural waters for trace metal analysis. *Introductory Journal of Environment* 11: 745-746.
- Boadi, K O and Kuitunen, M (2003). Municipal solid waste management in the Accra metropolitan area, Ghana. *The Environmentalist* 23:211-218.
- Bonde, G J (1977). "Bacterial indication of water pollution" In: *Advances in Aquatic Microbiology*, Academic Press, London
- Burubai, W; Akor, A J and Lilly, M T (2007). Performance evaluation of a septic system for high water-table areas. *American Eurasian Journal of Scientific Research* 2(2): 112116
- Bicki, T (2001). Onsite sewage disposal: the influence of system density on water quality. *Journal of Environmental Health* 53(5):39-42
- Chapman, D (1996). "Water Quality Assessment. A Guide to the Use of Biota, Sediments and Water in Environmental Monitoring," 2nd Edition, E&FN Spon, New York.



**Original Article**

- Chuku, O; Mustapha, H I and Abdul Gafar, H B (2008). The effect of Minna abattoir waste on surface water quality. *Environmental Research Journal* 2(6): 339-342
- Collins C H and Lyne M P (1976). *Microbiological Methods*. Butterworth and Co Publishers Ltd., London, Boston. 524.
- Collins, C H and Lyne, P M (1980). *Microbiological Methods*. Butterworth and Co.Limited, London. pp. 125-300.
- Coyne M S and Howell J M (1994). The faecal coliform/faecal streptococci ratio (FC/FS) and water quality in the Bluegrass Region of Kentucky. *Soil Science News and Views* 15. Daniels, R and Weaver, A (1987). Water table levels in some North Carolina soils. USDA Soils Conservation Service No. 603. Government Printing Office Washington, DC
- Ekugo,E I (1998). Public Health and Urban Sanitation. *Environ. News* 5, 7
- EPA (2002). “Ghana Landfills Guidelines”, Accra, 2002.
- Feng P; Weagant S and Grant M, (2007). Enumeration of *Escherichia coli* and the coliform bacteria. *Bacteriological Analytical Manual*. FDA/Center for Food Safety and Applied Nutrition.
- Geldreich, E E (1972). Buffalo lake recreational water quality: a study in bacteriological data interpretation. *Water Res. Pergamon Press*. 6:913-924.
- Gelt, J (1998). Microbes increasingly viewed as water quality threat. *Arroyo*. 10:1.
- Lateef, I, Manzoor A, Thoker, and Yousuf, A R (2003). Bacteriological survey of 15 springs of Kashmir. *Journal Research Development*. 3.
- Nartey, V K; Hayford, E K and Ametsi, S K (2012). Assessment of the impact of solid waste dumpsites on some surface water systems in the Accra Metropolitan Area, Ghana. *Journal of Water Resource and Protection* 4:605-61
- Ochuko, U and Thaddeus, O (2013). Effect of underground on-site sewage disposal system on the quality of water from hand dug wells in the urban centre of Ughelli, Delta State Nigeria. *Standard Journal of Education and Essay* 1(6): 81-90
- Ogbonna, D. N (2014). Distribution of Microorganisms in Water, Soils and Sediment from Abattoir Wastes in Southern Nigeria. *International Journal of Current Microbiology and Applied Science* 3(9): 1183-1200
- Oribaho, B J (2016). “Impact of human activities on biodiversity in Nigerian aquatic ecosystems” *Science International*. 4: 12-20
- Prescott, L M; Harley, J P and Klein, D A (2005). *Microbiology*. 6<sup>th</sup> ed. McGraw Hill, London. Pp. 135-140.
- Sadat, A, Akaki, K D; Ngoran, E B Z; Parinet, B and Frere J (2011). Evaluation of

**Original Article**

- Bacteriological Pollution of Yamoussoukro Lakes (Côte D'ivoire). *Canadian Research Journal of Biological Sciences*. 3(4):318-321.
- Seiyaboh E.I, Inyang I.R and Gijo A.H (2013). "Environmental impact of Tombia Bridge construction across Nun River in Central Niger Delta, Nigeria". *International Journal of Engineering and Science*. 2(11): 32-41
- Shafi, S, Kamili, A N; Shah, M A and Bandh, S A (2013). Coliform bacterial estimation: A tool for assessing water quality of Manasbal Lake of Kashmir, Himalaya. *African Journal of Microbiological Research* 7(31): 3996-4000
- Sharma, P; Sood, A, Sharma, S; Bisht, S; Kumar, V; Pandey, P; Gusain, M P and Gusain, O P (2010). Bacterial indicators of faecal pollution and physiochemical assessment of important North Indian lakes. *RMZ-Mater. Geoenvironment*. 57(1):25-40.
- Spellman, F R and Drinan, J (2000). "The Drinking Water Handbook," Technomic Publishing Co. Inc., Lancaster.
- Stewart, E (2005). Evaluation of septic tank and subsurface flow wetland for Jamaican Public school wastewater treatment. *M. Sc Thesis* Michigan Technological University.
- 64pp
- United Nations Environmental Programme, UNEP (2005). "Freshwater Pollution", *UNEP/GEMS Environmental Library*. 5(6):5-10
- United Nations International Emergency Children's Fund, UNICEF (1999). "Project Report: Study on Possible Pollution Caused by Water Seal Latrine to Ground Soil and Groundwater after 10 Years of Operation CEC," University of Science, Hanoi National University, Vietnam,
- United States Environmental Protection Agency (USEPA) (2009). "Innovative Uses of Composts," [www.epa.gov](http://www.epa.gov)
- Van, P and Pur, A (1990). The importance of clean water to Industries in Developed World 4<sup>th</sup> ed. Hong Kong, Grovener Press
- World Health Organization, WHO (2003). Guidelines for Safe Recreational-water Environments, Coastal and Fresh-waters Vol. 1. World Health Organization, Geneva, Switzerland
- World Health Organization, WHO (2009). "Guidelines for Drinking-Water Quality, Volume 1:Recommendations.Geneva:WHO,"[http://www.who.int/water\\_sanitation\\_health/dwq/GDWQ2004web.pdf](http://www.who.int/water_sanitation_health/dwq/GDWQ2004web.pdf)